

Journal of Chemical, Biological and Physical Sciences



An International Peer Review E-3 Journal of Sciences

Available online at www.jcbps.org

Section A: Chemical Sciences

CODEN (USA): JCBPAT

Research Article

Lead (II) and Chromium (IV) Biosorption from Aqueous Solutions by Green Algae Spirogyra Sp.

N. Gupta¹, J. K. Shrivastava², A.K. Gupta³

¹Department of Chemical Engineering, Ujjain Engineering College Ujjain 456010

²Department Civil Engineering, Ujjain Engineering College Ujjain 456010, India

Received: 08 January 2016; **Revised:** 01 February 2016; **Accepted:** 05 February 2016

Abstract: Heavy metal ion removal from aqueous solution or from wastewater by Biosorption Method is very cost effective and efficient Process. Results of metal ion removal from aqueous solution by using spirogyra sp. is presented here. Effect of pH, Time, Biosorbent Dose on the biosorption of heavy metals Pb(II) and Cr (VI) also described in this paper. Batch experiment were conducted to determine the biosorption capacity of spirogyra sp. and found results of maximum capacity of spirogyra sp. for lead is found to be 82% at pH 5 and at pH 2 for Cr is 75% with algal dose of 5 gram at 25^oC for both heavy metal ions. The effective pH Value for Pb (II) and Cr(VI) are found to be 5 and 5.5 , time required for metal uptake from wastewater was investigated for both metal ion and results found to be 90 minutes for Cr(VI) removal and 50 minutes for Pb(II) metal ion removal. Results shows that green algae spirogyra sp. is suitable biosorbent for the removal and with good recovery of Pb(II) and Cr(VI) from aqueous solution.

Keywords: Spirogyra sp., Algae, Heavy metals, biosorption.

INTRODUCTION

Water pollution due to the heavy metals is very serious problem in the world. The quality of life on earth is inextricably linked to the overall quality of the environment. The indiscriminate release of hazardous

pollutants by industries and dumping of domestic sewage in the human environment pose a major threat to all kinds of organisms inhabiting aquatic as well as terrestrial ecosystems¹. A heavy metal is a collective term for metals of high atomic mass, particularly those that are toxic and cannot be processed by living organisms. These include lead, chromium among others. Many other definitions of heavy metals have been proposed based on density, atomic number and atomic weight. Heavy metals have thus been defined collectively as metals of high atomic mass, particularly those transition metals that are toxic and cannot be processed by living organisms².

Lead and Chromium are among the most toxic heavy metals ion affecting the environment³. Lead comes into water through the combustion of fossil fuels and the smelting of sulphide ore, and into lakes and streams by acid mine drainage. Process industries, such as battery manufacturing and metal plating and finishing are also prime source of Pb pollution.

The current EPA and WHO drinking water standard for lead is 0.05 mg/L and 10 g/L, respectively. Lead accumulates mainly in bones, brain, kidney and muscles and may cause many serious disorders like anemia, kidney diseases, nervous disorders and sickness even death. It is therefore, essential to remove Pb(II) from wastewater before disposal. Chromium is one of the contaminants, which exists in hexavalent and trivalent forms. Hexavalent form is more toxic⁴ than trivalent and requires more concern. Strong exposure of Cr(VI) causes cancer in digestive tract and lungs⁵ and may cause epigastric pain, nausea, vomiting, severe diarrhea and hemorrhage⁶. It is therefore, essential to remove Cr(VI) from wastewater before disposal. The main sources of chromium (VI) are tannery, paint, ink, dye, and aluminum manufacturing industries etc.

Many treatment methods has been applied for the removal of heavy metals from waste water these are chemical precipitation, membrane filtration, ion exchange, reverse osmosis, electro dialysis, solvent extraction, evaporation, oxidation, activated carbon adsorption⁷. However, these methods are often expensive and difficult to maintain due to high capital and operational costs, as well as, extra cost of treating the resultant sludge generated before disposal⁸. So it is require to find best and secure technology for the removal of these toxic metals ions from waste water and I choose biosorption process for the treatment of waste water because the cost of it is less and high metal binding capacity. Aim of this study was to investigate that whether green algae *spirogyra* sp. Is suitable for the removal of lead and chromium ions from waste water (and under which experimental conditions: pH, contact time, *etc.*) AAS (Atomic Adsorption Spectrometry) was used for the determination of the Concentration of heavy metals in treated model solutions.

MATERIALS AND METHODS

2.1 Preparation of Biosorbent: The algae biomass was collected from lake near of Indore India. Before used it was wash from distilled water to remove the impurities and unwanted materials after washing it was kept on filter paper to reduce the water. The algae biomass was then dried in oven at 50⁰c for 12 hrs. The final weight was recorded and then crushed through a 250nm sieve to obtain a uniform particle size.

2.2 Glassware and Apparatus: All the glasswares (Beaker, Pipette, Conical flasks, volumetric flasks, Test tubes etc.) used were of Borosil/Rankem make. Before used, the apparatus washed with distilled water and then dried in oven at 65⁰C for 2 hrs.

2.3 Equipment: The equipment used throughout the experiment are listed in table below:

Table 2.1: List of Instrument used during the whole Experiment

S.No.	Instrument	Make
1.	Atomic Adsorption Spectrophotometer	GBC AVANTA PM HG3000
2.	Digital Weight Balance	Atco company
3.	What man filter Paper no. 1	-
4.	Orbital Incubator shaker	REMI Instrument, Mumbai
5.	pH meter	Adair dutt (AOP-Series)
6.	Hot air oven 142	Remi Instrument, Mumbai
7.	Magnetic hot plate stirrer	Neolab Company, Mumbai

2.4 Reagent used: All chemicals used in this work were of either analytical grade or Laboratory reagent (LR) grade obtained either from Merck, Germany or SD Fine Chem. Ltd India. Stock solution of lead and chromium were prepared by using lead nitrate and potassium dichromate in distilled water. Pb(II) and Cr (VI) solutions of different concentrations were obtained by diluting the stock solution. Standard solution of Pb(II) and Cr(VI) (1000 mg/L) for atomic adsorption spectrophotometer was obtained from Merck, Germany. Standard acid and base solutions (0.1N HCl and 0.1N NaOH) were used for pH adjustments.

2.5 Batch Adsorption studies: The biosorption of heavy metals on biosorbent was studied by batch technique. The adsorption features of the biosorbent *Spirogyra* sp. were investigated as a function of initial pH, initial heavy metal concentration, biosorbent dose and contact time.

The equilibrium and kinetics were obtained from batch experiments, using 250mL flasks containing 100mL of heavy metal solutions and 0.05 g of biomass kept at a fixed temperature (25°C). 1000mg/l stock solution of each of the metals was prepared, from where the working solutions were prepared by serial dilution method. The concentrations in mg/l for all the metals are 20mg/l. The pH value was adjusted to the required value with 0.1M HCl or 0.1N NaOH throughout the experiment. The remaining concentration of Pb(II) and Cr(VI) in residual solution was analyzed by taking absorbance on the atomic absorption spectrophotometer.

RESULTS AND DISCUSSION

3.1 Effect of contact time, pH and biosorbent dose on biosorption of Pb and Cr ions by green algae *Spirogyra* Sp.:

3.1.1 Effect of contact time Variation: The maximum uptake capacity of green algae *Spirogyra* sp. for Pb ion is 78.4% at 50 minute and for Cr ion 48.4% at 90 minute. Pb ions uptake increases up to by 20 minute of contact time, followed by a relatively slower rate up to 50 minutes and there after no significant uptake of Pb ion was observed (**Figure: 3.1**). For Cr ion equilibrium concentration was achieved with in 90 minute (at constant temperature 25°C). For Pb ion equilibrium concentration was achieved with in 50 minute (at constant temperature 25°C).




Figure: 3.1: Effect of time for removal of Pb and Cr ions by green algae *spirogyra* sp.

3.1.2 Effect of pH: pH is one of the important parameters in metal sorption by *spirogyra* sp. therefore metal sorption studies were carried out at different pH values. The biosorption of Pb and Cr ions were studied at different pH values, such as 2.0, 3.0, 4.0, 5.0, 6.0 and 7.0 (at constant temperature 25°C, Eq. time 50 minutes for Pb and 90 minutes for Cr ions). Optimum pH value observed 5 for Pb at and 5.5 for Cr Biosorption in metal ion solution as shown in **Figure 3.2**.

For all the metals considered, metal removal by all processes was low at low pH. For biosorption the removal rises to a peak between pH 5 and 5.5 then starts to decline. At low pH there is high competition for adsorption sites between metal ions and protons. Furthermore, at low pH both the hydrogen ion and the metal ion concentration are high since the metals are not precipitated but are available in solution. This explains why both processes are ineffective at low pH values. Since algal biomass has a high content of carboxyl groups on its cell walls, biosorption process can be affected by changes in the solution pH⁹. Interaction of dissociation sites on the biosorbent surface and sorbate solution chemistry depends upon the pH of the solution, such as hydrolysis, complexation by ligands, precipitation and availability of Pb and Cr ions, etc.




Figure: 3.2: Effect of pH for the removal of Pb and Cr ions by *spirogyra* sp.

3.1.3 Effect of Biosorbent Dose: To study the biosorbent dose on metal uptake, different amount of dose (0.5-5g) were used in 10ml solution of Pb(II) and Cr(VI) in which intake concentration of lead and Cr was 20mg/l. The amount of dose significantly affects the removal of Pb and Cr at optimum pH 5 for lead and at optimum pH 5 for chromium as shown in **Figure 3.3**. The percentage removal of lead was 32% at algal dose of 0.5 gram and it increased from 82% at algal dose of 5 gram. However there is slight changed when algal dose between 3-5 gram. The result found further increase in adsorbent dose, the capacity of adsorption found decrease. At low algal dose, all types of sites are entirely exposed and the adsorption on the surface is saturated faster. The percent removal of Cr was 8% at algal dose of 0.5 gram with contact time of 90 minutes and it increased to 75% at algae dose of 5 gram. It can is shown in figure 3.3 that there is slight change between from algal dose of 3-5 gram. This is consistent with the results obtained for the other adsorbent system¹⁰.




Figure 3.3: Effect of Biosorbent dose for the removal of Pb and Cr ions (20mg/l)

CONCLUSION

The present work indicates green algae *spirogyra* sp. is effective for the study is conduct to know the biosorption of heavy metal ions Pb(II) and Cr(VI) from aqueous solution. The optimum pH on the removal of Pb(II) and Cr(VI) found to be 5 and 5.5, the optimum contact time is 50minute for Pb(II) and 90 minutes for Cr(VI) ions. The maximum removal of lead is identify 82% at algal dose of 5 gram and the maximum removal of chromium is 75% from aqueous solution at algal dose of 5 gram. With the advantage of high metal biosorption capacity, the biomass of *Spirogyra* has the potential to be used as an efficient and economic biosorbent material for the removal of lead and chromium from aqueous solutions. From this work, green algae was found to be a biosorbent which can be used for effectively removing heavy metals from polluted water.

ACKNOWLEDGEMENTS

Authors are thankful to **Pollution Control Board Indore** for supporting this work.

REFERENCE

1. L. F. D. Filippis and C. K. Pallaghy, Heavy metals: Sources and biological effects. *Algae and Water Pollution*, 1994, 31-77.
2. P. Harrison and G. Waite, *The Cassell Dictionary of Chemistry*, Cassell, London, 1998.
3. V. K. Gupta, A. K. Srivastava and N. Jain, Biosorption of chromium (VI) from aqueous solutions by green algae *Spirogyra* species. *Water Res.*, 2001, 35, 4079–4085.
4. R. G. Smith and D. H. K. L. Lec, *Chromium in Metallic Contaminants and Human Health*, Academic Press, New York, 1972.
5. D.B. Kuafaman, Acute potassium dichromate poisoning in man. *AM. J. Diseases Children* 1970, 119, 374-379.
6. E. Browning, *Toxicity of Industrial Metals*, Chapter 12, 2nd ed. Butterworths, London, UK, 1969.
7. L. Pappalardo, F. Jumean, N. Abdo, “Removal of cadmium, copper, lead and nickel from aqueous solution by White, Yellow, and Red United Arab Emirates sand”, *American Journal of Environmental sciences*, 2010, 6(1), 41-44.
8. U. Kumar, “Agricultural Products and Byproducts As a Low-cost Adsorbent for Heavy Metal Removal from Water and Wastewater,” A Review, *Scientific Research and Essay*, 2006, 1(2), 033-037.
9. J.T Matheickal and Q. Yu, Biosorption of lead(II) and copper(II) from aqueous solutions by pre-treated biomass of Australian marine algae, *Bioresource Technology*, 1999, 69(3), 223-229.
10. I. Tuzun, G. Bayramoglu, Y.E. Alcin, G. Basaran, G. Celik, M.Y. Arica, Equilibrium and kinetic studies on biosorption of Hg(II), Cd(II) and Pb(II) ions onto microalgae *Chlamydomonas reinhardtii*, *J. Environ. Manage.* 2005, 77, 85-92.

*** Corresponding author: N. Gupta;**

Department of Chemical Engineering, Ujjain Engineering College
Ujjain 456010