

Journal of Chemical, Biological and Physical Sciences



An International Peer Review E-3 Journal of Sciences

Available online at www.jcbsc.org

Section C: Physical Sciences

CODEN (USA): JCBPAT

Research article

A Theoretical Study of two Dimensional Liquid ^4He and Evaluation of Ground State Energy as a Function of Densities Using two Potentials

Dinesh Kumar¹ and Lalit Kumar Mishra²

¹Kalawati Niwas, Kishori Mohan Complex, Road No 4, Po- Chandauti, Gaya-823001(Bihar) India

²Department of Physics, Magadh University (Bodh -Gaya) 824234(Bihar) India

Received: 5 October 2013; Revised: 26 October 2013; Accepted: 6 November 2013

Abstract: The ground state energy of two-dimensional liquid He4 was calculated as a function of densities using two potentials namely Lenard-Jones potential and Aziz potential. We have taken help of the theoretical formalism of Chung-In-Um et al (JLTP, 1997) in this evaluation. On comparing our theoretically evaluated results with GFMC calculation, it was observed that results obtained in the case of Aziz potential is very close to GFMC calculation.

Keywords : Learnd –Jones potential, Aziz potential, GFMC calculation, HNC and FHNC theories, radial distribution function, Liquid structure function, Three-body correlation function, Variation Monte Carlo simulation

INTRODUCTION

For many years, physicist have used hyper netted (HNC) and Fermi hyper netted chain (FHNC) theories to investigate the properties of highly correlated interacting boson and fermion systems such as liquid ^4He and normal liquid ^3He ^{1, 2}. The HNC/FHNC equations with Jastrow ground-state wave functions treat consistently both long and short correlations of the system and give agreements to some degree with experimental results. In spite of qualitative successes of the Jastrow ground-state function, there exist quantitative differences between HNC/FHNC results and experimental ones,

especially the equilibrium ground-state energy, and this fact leads one to consider the contributions arising from the three-body correlation functions³. It is well known that the three-body correlations do not affect seriously the structural properties of ground states, such as the radial distribution function and liquid structure function, but contribute significantly to the ground-state energy. Since liquid ³He system is less dense than liquid ⁴He, that is, the equilibrium density of the former is 0.0166 Å⁻¹ while that of the latter is 0.02185 Å⁻³ in a three-dimensional system, effects of three-body correlations in ³He are rather small compared to those in ⁴He. At equilibrium density, they make up about half the difference between the Jastrow results and experimental energy in a liquid ³He system⁵ and lower the Jastrow ground-state energy by about 10 % in liquid ³He.

Trial ground-state wave function including three-body correlation functions in addition to two-body correlation effects can be constructed in the form

$$\psi(\bar{r}_1, \bar{r}_2, \dots, \bar{r}_N) = \prod_{i < j}^N f_2(r_{ij}) \prod_{i < j < k}^N f_3(r_{ij}, r_{jk}, r_{ki}) |\Phi\rangle = F|\Phi\rangle$$

Where $f_2(r_{ij})$ is the two-body correlation function which describes spatial correlations and depends only upon the spatial distance r_{ij} between two particle $f_3(r_{ij}, r_{jk}, r_{ki})$ is a three-body correlation function, and $|\Phi\rangle$ is the anti-symmetric product of non-interacting single-particle plane waves normalized in the system's area Ω with N fermions, which becomes simply unity for boson systems.

In addition to the three-body correlation functions in the ground-state wave function, one should take into account the contributions coming from the elementary diagrams as well as effects of the HNC/FHNC equations to obtain better results. Unfortunately, there does not exist a general closed expression for the elementary diagrams and therefore one must calculate each diagram individually, which is very time consuming. During the last two decades, two approximations for the estimation of the elementary diagrams have been developed. One is known as the scaling approximation⁴ and the other is the interpolating equation approximation⁵. Both techniques have shown their efficiency for central two-body correlations, providing similar results for the total ground-state energy to those of the variational Monte Carlo simulation.

The scaling approximation uses the fact that successive contributions of the five, six and higher order elementary diagrams are approximately proportional to and have very similar spatial behavior to the body elementary diagrams⁴, so that one can represent the total elementary diagram contribution by means of the readily calculable four-body elementary diagrams. The interpolating equation approximation determines a parameter such that it gives the same value for the classical isothermal compressibility obtained from both the HNC/FHNC equations and Percus-Yevick⁶ equations using the classical pressure derivative or the compressibility integral.

Although large number of theoretical works regarding ⁴He and ³He system have been reported⁷⁻¹¹ but very few of them have been used for two-dimensional liquid.

Chung-In Um *et al.*¹² have developed a theoretical model to study two-dimensional quantum fluid. By taking their model, one has evaluated the radial distribution function $g(r)$ as a function of r for two dimensional liquid ⁴He interacting through Aziz potential for various densities. In **section 2**, we have given the mathematical formulae used in the evaluation. The discussion of the result has been given in **section 3** of this paper. In the last portion, we have shown the tables of the results and the reference taken.

MATHEMATICAL FORMULAE USED IN THE EVALUATION

The Hamiltonian for the considered system is given by

$$H = T + V = \sum_i T_i + \sum_{i < j} V_{ij} = -\frac{\hbar^2}{m} \sum_i \nabla_i^2 + \sum_{i < j} V(\vec{r}_i - \vec{r}_j) \quad (1)$$

Where $V(r)$ is the interatomic interaction. A popular potential theoretical investigations is that of Lenard-Jones, which has parameters, i.e., the hard-core radius and the well depth, and can be written as

$$V_{LJ}(r) = 4\epsilon \left[\frac{12}{r_m^6} - \frac{1}{r_m^{12}} \right] = 1022 K, \sigma = 2.556 \text{ \AA} \quad (2)$$

However, $V_{LJ}(r)$ includes only the dipole-dipole interaction and does not take into account multiple interactions. A more physically realistic potential which accounts for the self-consistent field Hartree-Fock repulsion and multiple interactions is the HFDHE² potential of Aziz *et al.*¹³ presented as

$$V_{LJ}(r) = \epsilon^* \exp\left(-\alpha \frac{r}{r_m}\right) \left[c_6 \frac{1}{r_m^6} + c_8 \frac{1}{r_m^8} + c_{10} \frac{1}{r_m^{10}} \right] F(r) \quad (3)$$

Where

$$F(r) = \begin{cases} \exp[-(Dr_m/r - 1)^2] & r/r_m \leq D \\ 1 & r/r_m \geq D \end{cases} \quad (4)$$

The values of the constants are

$$A = 0.54485 \times 106, \quad \epsilon^*/k_B = 10.8$$

$$\alpha = 13.353384, \quad C_6 = 1.3732412$$

$$C_8 = 0.4253785, \quad C_{10} = 0.178100$$

$$D = 1.241314, \quad r_m = 2.9673 \text{ \AA}$$

It is well known from the Green's function Monte Carlo simulations and other variational calculations that the Aziz potential gives closer results to experiments than V_{LJ} in three dimensions. One uses both potentials and compare the results with those in two dimensions.

The n-particle distribution function is represented as

$$P_n(\vec{r}_1, \dots, \vec{r}_n) = v^n N(N-1) \times \frac{\int \psi^*(\vec{r}_1, \dots, \vec{r}_n) \psi(\vec{r}_1, \dots, \vec{r}_n) d\vec{r}_{n+1} \dots d\vec{r}_N}{\langle \psi | \psi \rangle} \quad (5)$$

Where \vec{r}_i denotes the spatial coordinates of the i^{th} particle and v is the spin degeneracy of the system (2 for this system). In homogeneous system, as $N \rightarrow \infty$ and $\Omega \rightarrow \infty$, the single-particle distribution function reduces to the density of the system so that the radial distribution function can be expressed in terms of the density and two-particle distribution function $P_2(r)$ as

$$g(r) = \frac{1}{\rho^2} p_2(r) = N(N-1) \frac{\int_{\rho}^{k_F} \sum_{\psi^+}(\vec{r}_1, \dots, \vec{r}_n) \psi(\vec{r}_1, \dots, \vec{r}_n) d\vec{r}_3 \dots d\vec{r}_N}{\langle \psi | \psi \rangle} \quad (5)$$

One uses the FHNC approximation to sum the diagrams arising from the cluster properties of g(r). In this scheme, g(r) can be decomposed as

$$g(r_{ij}) = g_{dd}(r_{ij}) + 2g_{de}(r_{ij}) + g_{ee}(r_{ij}) \quad (6)$$

Where dd, de and ee represent terms in which both i and j are not exchanged, only j is exchanged, and both i and j are exchanged, respectively. The components of g(r_{ij}) are given by

$$g_{dd}(r_{ij}) = \exp[u_2(r_{ij}) + N_{dd}(r_{ij}) + E_{dd}(r_{ij})]$$

$$g_{de}(r_{ij}) = g_{dd}(r_{ij})[N_{de}(r_{ij}) + E_{de}(r_{ij})]$$

$$g_{ee}(r_{ij}) = g_{dd}(r_{ij})[-L^2(r_{ij})/v + N_{ee}(r_{ij}) + E_{ee}(r_{ij}) + (N_{ee}(r_{ij}) + E_{ed}(r_{ij}))^2] \quad (7)$$

$$g_{ee}(r_{ij}) = g_{dd}(r_{ij})L(r_{ij})/v$$

Where

$$L(r_{ij}) = -l(k_F r_{ij}) + v(N_{ee}(r_{ij}) + E_{ee}(r_{ij})) \quad (8)$$

$$l(x) = 2j_1(x)/x \quad (9)$$

k_F is the Fermi momentum of the system, and $j_1(x)$ is the Bessel's function of the first kind of order 1. N_{mm} represent sums of the diagrams and E_{nm} sums of the elementary diagrams. The equation $g_{ee}(r_{ij})$ denotes the terms in which both i and j are exchanged in incomplete exchange loop, and $N_{ee}(r_{ij})$ and $E_{ee}(r_{ij})$ are sums of the nodal diagrams and the elementary diagrams in which i and j belong to the same permutation loop, respectively.

With the FHNC equation and expression for the g(r), one now calculates the energy per particle of the system

$$\frac{E_{var}}{N} = \frac{1}{N} \frac{\langle \psi | H | \psi \rangle}{N \langle \psi | \psi \rangle} = \frac{\langle V \rangle}{N} + \frac{\langle T \rangle}{N} \quad (10)$$

The potential energy per particle is expressed by taking the thermodynamic limit, i.e., the area of system Ω and the number of particles N are infinite, keeping the density constant as

$$\frac{\langle V \rangle}{N} = \frac{\rho}{2} \sum_{ij} d\vec{r}_{ij} V(r_{ij}) g(r_{ij}) \quad (11)$$

The expression for the kinetic energy part is more complicated. There are three well-known expressions for the helium system i.e., Jackson-Feenberg¹⁴, Clark-Westhous¹⁵ and Bethe-Pandharipande¹⁶. Among expressions for the kinetic energy per particle the Jackson-Feenberg form is popular and easily tractable since its three-particle distribution part is not more significant than others. One uses here the Jackson-Feenberg and the Bethe-Pandharipande forms so that we make use of the scaling approximation for the elementary diagrams. From the identity

$$\nabla^{\dagger} \nabla^2 \psi d\vec{r} = \frac{1}{4} \nabla \psi^{\dagger} (\nabla^2 \psi) (\nabla^2 \psi^{\dagger}) + -2(\nabla \psi^{\dagger}).(\nabla \psi) d\vec{r}] \quad (12)$$

Jackson-Feenberg energy form is obtained as

$$\begin{aligned} \frac{E_{JF}}{N} &= \frac{-\hbar^2}{2mN} \frac{1}{\langle \psi | \psi \rangle} \sum_i \left\langle \Phi | F^2 \nabla^2 \nabla_i^2 + \frac{1}{2} F \nabla_i^2 F - \frac{1}{2} (\nabla_i F)^2 | \Phi \right\rangle \\ &+ \frac{\hbar^2}{2mN} \frac{1}{\langle \psi | \psi \rangle} \sum_i F^2 \nabla_i^2 |\Phi|^2 d\vec{r}_1 \dots d\vec{r}_N + \frac{\langle V \rangle}{N} \\ &= T_F = W_B + W_\phi + U_\phi \end{aligned} \quad (13)$$

The first term of Eq. (13), $TF = \frac{\hbar^2 k_F^2}{4m}$, is the non interacting fermion energy with the Fermi momentum $k_F = [4\pi^2 N / V]^{1/3}$, and the second, WB is the energy of the boson system.

$$W_B = \frac{\rho}{2} \nabla^2 r g(r) - \frac{\hbar^2}{(2m)} \frac{\nabla^2 f_2(r) - (\nabla f_2(r))^2}{f_2(r)} \quad (14)$$

The forms of W_ϕ and U_ϕ are

$$\begin{aligned} W_\phi &= \frac{\hbar^2 \rho}{4mv} \nabla^2 r \{ g_{dd}(r) - 1 \} (k_F r) - v (N_{ee}(r) + E_{ee}(r)) \nabla^2 l(k_F r) \\ &\times \nabla^2 l(k_F r) + \nabla l(k_F r)^2 - v d_{dd}(r) E_{cc}(r) \nabla^2 l(k_F r) \} \end{aligned} \quad (15)$$

$$\begin{aligned} U_\phi &= \frac{\hbar^2 \rho^2}{8m} \nabla^2 r_{12} d^2 r_{13} (g_{dd}(r_{12}) - 1) \nabla l(k_F r_{12}) \cdot (g_{dd}(r_{13}) - 1) \\ &\times \nabla l(k_F r_{13}) g_{ee}(r_{23}) \end{aligned} \quad (16)$$

$$U_\tau = \frac{\hbar^2 \rho}{16m} \nabla^2 r_{12} d^2 r_{13} g_3(r_{12}, r_{23}, r_{13}) \nabla^2 u_3(r_{12}, r_{23}, r_{13}) \quad (17)$$

Pandharipande-Bethe energy form is obtained by expanding the kinetic energy per particle as

$$-\frac{\hbar^2}{2mN} \frac{1}{\langle \psi | \psi \rangle} \sum_i \left\langle \Phi | F \nabla_i^2 [F] | \Phi \right\rangle = \frac{\langle T \rangle_{PB}}{N} \quad (18)$$

From which one gets,

$$\frac{\langle T \rangle_{PB}}{N} = -\frac{\hbar^2}{2mN} \frac{1}{\langle \psi | \psi \rangle} \sum_i \left\langle \Phi | F [\nabla_i^2 + \nabla_i^2 + 2(\nabla_i F)] | \Phi \right\rangle \quad (19)$$

After some calculations, we obtained E_{PB}/N as

$$\frac{E_{PB}}{N} = T_F + W + W_F + U + U_F \quad (20)$$

Where W and U are the bosonic energies per particle, and W_F and U_F are from the last term of Eq. (19)

$$W = \frac{\rho}{2} \sum r_g(r) \left[-\frac{\hbar^2}{m} \nabla^2 f_2(r)^2 \right] \quad (21)$$

$$U = -\frac{\hbar^2 \rho^2}{2m} \sum r_{12} d^2 r_{13} g_3(r_{12}, r_{23}, r_{13}) \left[\frac{\nabla f_2(r_{12}) \cdot \nabla f_2(r_{13})}{f_2(r_{12}) f_2(r_{13})} \right] \quad (22)$$

$$W_F = -\frac{\hbar^2 \rho}{2m} \sum r g_{ee}(r) \left[\frac{\nabla f_2(r) \cdot \nabla l(k_F r)}{f_2(r)} \right] \quad (23)$$

$$U_F = -\frac{\hbar^2 \rho^2}{2m} \sum r_{12} \nabla f_2(r_{12}) \cdot \nabla l(k_F r_{13}) \\ \times \{g_{ee}(r_{13}) g_{dy}(r_{12})(g_{dy}(r_{32}) - 1) + g_{dd}(r_{13}) g_{ee}(r_{23}) g_{ee}(r_{31})\} \quad (24)$$

For numerical calculations, one adopts the McMillan type function as a trial two-body correlation function, which is widely used in variational and Monte Carlo calculations for liquid and helium systems.

$$u_2(r) = -\frac{b}{r} \quad (25)$$

Where, b is a variational parameter chosen to give an optimized distribution function at a given density. To obtain more precise results, we include three-body correlation effects in the following forms:

$$u_3(r_{ij}, r_{jk}, r_{ki}) = \sum_{cyc} \eta(r_{ij}) \eta(r_{jk}) (\hat{r}_{ij} \cdot \hat{r}_{jk}) \quad (26)$$

$$\eta(r) = \sqrt{\lambda_1 r} \exp \left[-\frac{r}{b_1} \right] \quad (27)$$

Where cyc denotes the cyclic permutation among the three particles coordinates i, j and k, and \hat{r}_{ij} is a unit vector along the line connecting particle i and j. Since it was first used in a variational Monte Carlo calculation⁸. This form for the three-body correlations is generally used in variational HNC/FHN methods¹⁻³.

DISCUSSION OF RESULTS

In this paper, we have evaluated the ground state energy of two dimensional liquid ${}^4\text{He}$ interacting through Lenard-Jones and Aziz potential at various densities. We have used the theoretical formalism of Chung-In Um *et al.*¹² in our evaluation. Chung-In Um¹² calculated ground state energy of two dimensional liquid ${}^4\text{He}$ using Jastrow two body correlation function in the trial wave function within the HNC approximation. His other calculation for energy per particle is by taking two body correlation and contribution of order elementary diagrams. We have taken three body as well as two -

body correlations within the scaling approximation⁴. Our theoretically evaluated results of ground state energy E(k)/N are shown in **Table T1** and **T2** for Lenard –Jones potential and Aziz potential for various densities. We have compared our theoretical results with Chang's result¹⁷ and GFMC result⁸. Our theoretical calculation indicates that the two dimensional liquid ⁴He interacting through Aziz potential is more close to the calculation⁸. Comparing with two potential, one conclude that Aziz potential is more suitable to describe, the ground state of the two dimensional helium system similar to bulk case 18. Some recent¹⁹⁻²⁵ results reveal the same fact.

Table T1: Evaluated results of ground state energy of two-dimensional liquid ⁴He interacting through the Lenard-Jones potential at various densities.

$\rho(\text{A}^{\circ -2})$	Ground state energy E(k)/N		
	Our cal. results	Chang's ¹⁷ results	GFMC Calculations ⁸
0.020	-0.520	-0.628	-0.504
0.025	-0.546	-0.637	-0.546
0.030	-0.624	-0.658	-0.589
0.035	-0.658	-0.586	-0.627
0.040	-0.748	-0.552	-0.553
0.045	-0.786	-0.505	-0.524
0.050	-0.842	-0.486	-0.508
0.055	-0.725	-0.453	-0.493
0.060	-0.556	-0.446	-0.486

Table T2: Evaluated results of ground state energy [E(k)/N] of two-dimensional liquid ⁴He interacting through Aziz potential at various densities

$\rho(\text{A}^{\circ -2})$	Ground state energy E(k)/N		
	Our cal. results	Chang's ¹⁷ results	GFMC Calculations ⁸
0.020	-0.487	-0.586	-0.523
0.025	-0.509	-0.603	-0.588
0.030	-0.586	-0.658	-0.602
0.035	-0.627	-0.709	-0.657
0.040	-0.723	-0.682	-0.597
0.045	-0.706	-0.596	-0.602
0.050	-0.658	-0.542	-0.524
0.055	-0.532	-0.506	-0.486
0.060	-0.456	-0.487	-0.442

REFERENCES

1. E. Krotscheck, *Phys. Rev.*(1986) **B 33**, 3158
2. F. Arias de Saavedra and E. Buendia, *Phys. Rev.*(1990) **B 42**, 6018
3. E. Manouskais, S. Fantoni, V.R. Pandharipande and Q.N Usmani, *Phys. Rev.*(1983) **B 28**, 3770
4. Q.N. Usmani, B. Friedman and V.R. Pandharipande, *Phys.Rev* (1982) **B 25**, 4502
5. A. Fabrocini and S. Rosati, Nuvino Cimento (1982) **D 1**, 567, 615
6. J.K. Percus and G.J. Yevick, *Phys. Rev.*(1958) 110, 1
7. K. S. Liu, M. H. Kalos and G. V. Chester ,*Phys. Rev*(1976).**B13**,1971
8. P. A. Whitlock, D. M. Ceperley, G. V. Chester and M. H. Kalos, *Phys. Rev.*(1979)**B19**,5598
9. P. Hatzikonstantinou, *J. Phys.*(1985) **C18**, 2393
10. P.A. Whitlock, G. V. Chester and M. H. Kalos, *Phys. Rev.*(1988) **B38**, 2418
11. S. Giorgini, J. Boronat and J. Casulleras, *Phys Rev.*(1996) **B54**,6099
12. Chung-In-Um, J. R. Kahng, Y. S.Kim, T. F. George and L. N. Pandey J. Low Tem. *Phys.*(JLTP)(1997) **107**, 283
13. R.A. Aziz, V.P.S. Nain, J.S. Carley, W.I. Taylor and G.T. Conville, *J. Chem. Phys.*(1979) **70**, 4330
14. H.W. Jackson and E. Feenberg, *Ann. Phys.* (N.Y.)(1961) **15**, 266
15. J.W. Clark and P. Westhaus, *Phys. Rev.* (1966) **141**,833
16. V.R. Pandharipande and H.A. Bethe, *Phys. Rev.* .(1973) **C 7**, 1312
17. C.C. Chang, J. Low. *Temp. Phys.* (JLTP)(1980) **38**, 25
18. M. Wagner and D. M. Ceperley J. Low. *Temp. Phys.*(JLTP)(1994) **94**,185
19. B. K. Clark and D. M. Ceperley, *Phys. Rev. Lett.*(PRL)(2006) **96**, 105302
20. C. Cazorla and J. Boronot , *Phys. Rev.*(2006) **B78**, 134509
21. C. Cazorla, D. Errandonea and E. Sola, *Phys. Rev.*(2009) **B80**, 064105
22. W.Kitterle, Y. Shin, A. Schivatzek and C. H. Shunk, *Int. J. Mod. Phys.*(2010) **B24**, 3950
23. F. Mallet, T. Kristensen, S. M. Reimann and G. M. Kavoulakin, *Phys. Rev.* (2011) **A83**, 033628
24. D. Blank, K. M. Daily and D. Rakshit, *Phys. Rev* (2012) **A85**, 021601
25. P. O. Bugunion, J. A. Lofthouse and G. J. Conduel, *Phys. Rev. Lett.* (PRL) 111, 045301

Corresponding author: Dinesh Kumar; Kalawati Niwas, Kishori Mohan Complex, Road No 4,
Po- Chandauti, Gaya-823001(Bihar)