

# Journal of Chemical, Biological and Physical Sciences



An International Peer Review E-3 Journal of Sciences

Available online at [www.jcbpsc.org](http://www.jcbpsc.org)

Section C: Physical Science

CODEN (USA): JCBPAT

Research article

## An Evaluation of Critical Temperature for Bose-Einstein Condensation of the Atom-Molecule Mixture at the Fixed Density

Indu Kumari Tiwari\*, R. N. Jha\*\* and L. K. Mishra

Department of Physics, Magadh University, Bodh-Gaya-824234 (Bihar)

\*\*Department of Physics, Maharaja College, Ara-802301, Bhojpur (Bihar)

\*c/o Shree Anant Kumar Tiwari, Village-Dhawari, P.o-Bara Khayawani, Dist-Bhojpur -802166 (Bihar)

Received: 20 September 2013; Revised: 10 October 2013; Accepted: 15 October 2013

**Abstract:** Using theoretical formalism of *R. A. Duine and H. T. C. Stoof (Physics Reports, 2004)* we have evaluated the critical temperature  $T_C$  for Bose-Einstein condensation (BEC) for the atom-molecule mixture at the fixed total density. The evaluation has been performed for positive and negative detuning. For positive detuning, we obtained  $T_{BEC} = T_0$  and for negative detuning, we have  $T_{BEC} = (2^{-5/3}T_0)$  where  $T_0$  is initial temperature. Our theoretically evaluated results are in good agreement with the experimental data and also with other theoretical workers.

**Keywords:** Real-time dynamics, Effective atom-molecular theory, Heisenberg equation, Retarded molecular propagator, Positive and negative detuning, Effective atom-molecular coupling, Wigner –threshold law, Backward scattering length.

### INTRODUCTION

In order to evaluate the critical temperature for Bose-Einstein condensation of the atom-molecule mixture at the fixed total atomic density, one studies the effective atom-molecular theory<sup>1,2</sup>. In this study, one considers the real-time dynamics of the system by deriving the Heisenberg equation of motion for the field operators that annihilate an atom and a molecule of a given space and time. To

determine the Heisenberg equation of motion for these field operators, one first have to perform an analytical continuation from Matsubara frequencies to real frequencies. For simplicity, one assumes that one is close to the resonance and one is allowed to neglect the energy dependence of the effective atomic interactions and the effective atom-molecule coupling. Moreover, one takes only the leading order energy dependence of the molecular self-energy into account.

Higher orders are straight forwardly included. The leading order energy dependence of the self energy is after Wick rotations and is real energies. For the positive energy E, this result is in agreement with Wigner- Threshold law<sup>3</sup>. This law gives the rate for a static with a well defined positive energy to decay into three dimensional continuum. In order to study the properties of the gas in the normal state, one considers the two- atom properties of many-body theory.

One can show that the effective field theory<sup>4</sup> correctly contains the two-atom physics of a Feshbach resonance<sup>5</sup>. Firstly, one shows that the correct Feshbach- resonant atomic scattering length is obtained after the elimination of the molecular field. Secondly, one calculates the bound energy and shows that it has the correct threshold behavior near the resonance. To get more insight in the nature of the molecular state near the resonance, one also investigates the molecular density of states.

To calculate the effective interatomic scattering length, one has to eliminate the molecular field from Heisenberg equation of motion. Since the scattering length is related to the scattering amplitude at zero energy and momentum, one is allowed to put all the time and spatial derivatives in the equation of motion for the molecular field operator equal to zero. The scattering length wear a Feshbach resonance and is given by

$$a(B) = a_{bg} + a_{res}(B)$$

where  $a_{bg}$  is the backward scattering length and  $a_{res}(B)$  is the Feshbach resonance. The energy of the molecular state is determined by the poles of the retarded molecular propagator. For positive detuning  $\delta(B)$ , there exists a pole with a non-zero and negative imaginary part. This is in agreement with the fact that a molecule decays when its energy is above the two atom continuum threshold. The imaginary part of the energy is related to the lifetime of the molecular state. For negative detuning the molecular propagator has a real and negative pole corresponding to the bound state energy.

The molecular density of state is obtained by taking the imaginary part of the retarded molecular propgator<sup>6</sup>. One can discuss only the situation when one is close to resonance and approximate the retarded molecular elf-energy by the square root term resulting from Wigner threshold law. For the case of negative detuning, the molecular density of states has two contributions, one arising from the pole at the bound state energy and the second from two atom continuum.

Then one defines the wave function renormalization factor Z(B). This factor goes to zero as one approach the resonance and it becomes equal to one far off resonance. Far off resonance, the bound state of the coupled channels i.e the dressed molecules is almost equal to the bound state of the closed channel potential and has zero amplitude in the open channel.

This corresponds to the situation  $Z(B) \sim 1.0$ . As the resonance is approached, the dressed molecule contains only with an amplitude  $\sqrt{Z(B)}$  the closed channel bound state i.e. bare molecule. Accordingly, the contribution of the open channel becomes larger and gives rise to the threshold behavior of the bound state energy. For positive detuning the molecular density of state has only a contribution for positive energy. The lifetime of the molecular state is defined by

$$\Gamma_m(B) = \frac{g^2 m^{\frac{3}{2}}}{\pi \hbar^4} \sqrt{\delta(B)}$$

$g$  is the order of coupling between atom and molecule,  $m$  is the mass of the atom,  $\delta(B) = \Delta\mu(B - B_0)$  is the detuning. As expected, the density of states is, in the case of positive detuning, approximately a Lorentzian expected around the detuning with a width related to the lifetime of the molecule<sup>7-10</sup>.

In this paper, we have evaluated the critical temperature for the Bose-Einstein condensation (BEC) for the atom-molecule mixture at the fixed total atomic density. Two results were given one for the exact calculation and other for treating the gas as an ideal gas mixture. The calculations were performed from positive to negative detuning.

**Mathematical formulae used in the evaluation:** The equilibrium properties of the gas are determined by the equation of state, which relates the total density of the gas to the chemical potential. The equation can be calculated in two ways, either by calculating the thermodynamic potential and differentiating with respect to the chemical potential, or by directly calculating the expectation value of the operator for the total density. First, one calculates the thermodynamic potential which is given by<sup>11</sup>

$$\Omega(\mu, T) = \frac{1}{\beta} T_r [\ln(G_{0,a}^{-1})] + \frac{1}{\beta} T_r [\ln(G_m^{-1})] \quad (1)$$

Here  $G_{0,a}(k, i\omega_n)$  is non interacting atomic propagator of the atoms.  $G_m(k, i\omega_n)$  is molecular propagator.  $\beta = (K_B T)^{-1}$ . The full molecular propagator is given by

$$G_m(k, i\omega_n) = \frac{-\hbar}{i\hbar\omega_n} + \frac{\varepsilon_k}{2} + \delta(b) - 2\mu + \hbar \sum_m^{2B} (i\hbar\omega_n - \frac{\varepsilon_k}{2} + 2\mu) \quad (2)$$

Here  $\hbar\Sigma_m^{2B}$  is the molecule self-energy. The total atomic density is calculated by using the thermodynamic identity  $N = \frac{-\partial\Omega(\mu, T)}{\partial\mu}$  which results in total density as

$$n = -\frac{1}{V} \sum_k [N(\varepsilon_k - \mu) + 2N(\frac{\varepsilon_k}{2} + \varepsilon_m(B) - 2\mu)] \quad (3)$$

Where  $\varepsilon_m(B)$  is molecular binding energy. This important results show that in equilibrium in the normal state and for negative detuning the gas in first approximation behaves as an ideal gas mixture of atoms and the dressed molecules. The same result is found if one neglects in the Heisenberg equations of motion for the atomic and dressed molecular field operators in the interaction terms and calculate the total density in equilibrium. Then, one calculates the total density for negative detuning the contribution comes from molecular propagator. It is found to be equal to<sup>12</sup>

$$n_m = Z(B) \int \frac{dk}{(2\pi)^3} \int N \left( \frac{\varepsilon_k}{2} + \varepsilon_m(B) - 2\mu \right) \quad (4)$$

Where  $Z(B)$  is wave function renormalization factor. The molecular self-energy is given by

$$\hbar\Sigma_s(k, i\omega_n) = \frac{4g^2}{V} \sum_{q,n} \frac{1}{[i\hbar\omega_n - (\delta(B)) - \varepsilon_k - \mu]} \quad (5)$$

Finally the total density of the atoms is given by

$$n_n = \frac{(2-2Z(B))}{V} \sum_k N \left( \frac{\varepsilon_k}{2} + \delta(B) - 2\mu \right) + \frac{1}{V} \sum_k N (\varepsilon_k - \mu) \quad (6)$$

The molecular density becomes equal to

$$n_m = \frac{Z(B)}{V} \sum_k N (\varepsilon_k + \delta(B) - 2\mu) \quad (7)$$

The temperature is given in units of the critical temperature for Bose-Einstein condensation of an ideal gas of atoms with total density n as

$$T_0 = \frac{3.31 \hbar^2 n^{2/3}}{m K_\beta} \quad (8)$$

**Adiabatic Sweep through the resonance:** One now calculates the number of atoms and molecules of the gas during adiabatic sweep in the magnetic field, such that the detuning changes from positive to negative<sup>13</sup>. The condition for adiabatically is that the entropy of the gas is constant. The entropy is given by

$$S = - \frac{\partial \Omega}{\partial T} \quad (9)$$

where  $\Omega$  is thermodynamic potential. The total number of atoms is also constant throughout the sweep. As one can see for sufficiently large absolute values of the detuning, the gas is well described by an ideal gas approximation. For simplicity, one will treat the gas here as an ideal-gas mixture since one is mostly interested in the final density of atoms and molecules and final temperature of the gas after the sweep<sup>14</sup>.

## DISCUSSION OF RESULTS

In this paper, we have evaluated the critical temperature for Bose-Einstein condensation (BEC) for the atom-molecule mixture at the fixed atomic density. The evaluation has been performed by the theoretical formalism of R. A. Duine and H. T. C. Stoof<sup>15</sup>. In **Table 1**, we have shown the temperature of the gas as a function of detuning for a sweep through the resonance from positive to negative detuning for two initial temperature  $T=2 T_0$  and  $4T_0$ . The total density of the gas is equal to  $n=10^{13} \text{ cm}^{-3}$ . The gas is heated as the detuning is changed from positive to negative. In **Table 2**, we have given the evaluated results of critical temperature for Bose- Einstein condensation (BEC) of atom-molecule mixture at a fixed total atomic density  $n=10^{13} \text{ cm}^{-3}$ .

Two results were given, one for the exact calculation and other for treating the gas as an ideal gas mixture. For positive detuning and far from the resonance, one is essentially dealing with an atomic gas. Hence one has in this regime  $T_{\text{BEC}} = T_0$ . For sufficiently negative detuning, one is dealing with a gas of molecules with twice the atomic traps and one has  $T_{\text{BEC}} = (2^{-5/3} T_0)$ . The feature in the critical temperature at zero detuning turns out to be a signature of true thermodynamic phase transition between a phase with single Bose-Einstein condensates of molecules and a phase contrary of two Bose-Einstein condensates, one of atoms and other of molecules. This was first pointed out by Sachher<sup>16</sup>. This should be contrasted with the situation of atomic Fermi gas near a Feshbach resonance, where only a BCS-BEC cross-over exists<sup>17</sup>.

In **Table 3**, we have shown the results of evaluation of fraction of atoms as a function of the detuning for an adiabatic sweep through the resonance. The total density is again kept equal to ( $n=10^{13} \text{ cm}^{-3}$ ). The calculation were performed with two initial temperature  $T=2T_0$  and  $4T_0$ . In **Table 4**, we have given the result of evaluation of twice the fraction of molecules as a function of detuning for an adiabatic sweep through the resonance. The other parameters are same as shown in **Table 3**. There are some recent results<sup>18-24</sup> which also reveal the similar behavior.

**Table 1:** Temperature of the gas as a function of the detuning for a sweep through resonance from positive to negative detuning for two initial temperature  $T=2T_0$  and  $4T_0$ . The total atomic density of the gas is  $n=10^{13} \text{ cm}^{-3}$ .

$\frac{4\pi^2\hbar^6\delta}{g^4m^3}$ Detuning	$T/T_0 (T=2T_0)$	$T/T_0 (T=4T_0)$
-0.0030	3.25	7.85
-0.0025	3.00	6.52
-0.0020	2.92	6.20
-0.0015	2.86	5.86
-0.0010	2.72	5.48
-0.0005	2.58	5.12
0.0	2.50	4.95
0.001	2.80	5.25
0.002	2.95	5.49
0.003	3.10	5.75
0.005	3.17	5.90

**Table 2:** An Evaluation of critical temperature for Bose-Einstein condensation as a function of detuning .The total density of the gas is equal to  $n=10^{13} \text{ cm}^{-3}$ . I- results show the exact calculations. II-results show the calculation for treating the gas as an ideal gas mixture.

$\frac{4\pi^2\hbar^6\delta}{g^4m^3}$ (detuning)	$( T_{BEC}/T_0 )$	
	result	result
-0.0030	0.345	0.347
-0.0025	0.348	0.354
-0.0020	0.352	0.359
-0.0015	0.358	0.365
-0.0010	0.362	0.369
-0.0005	0.371	0.375
0.00	0.385	0.352
0.001	0.486	0.364
0.002	0.685	0.458
0.003	0.795	0.886
0.004	0.928	0.958

**Table 3:** An evaluated results of function of atoms as a function of the detuning for an adiabatic sweep through the resonance. The total density is equal to  $n = 10^{13} \text{ cm}^{-3}$ . The calculations have been performed for two initial temperature  $T = 2T_0$  and  $4T_0$ .

$\frac{4\pi^2\hbar^6\delta}{g^4m^3}$	Fraction of atoms	
	$T = 2T_0$	$T = 4T_0$
-0.0030	0.00	0.252
-0.0025	0.005	0.284
-0.0020	0.008	0.327
-0.0015	0.015	0.368
-0.0010	0.225	0.445
-0.0005	0.325	0.565
0.00	0.458	0.589
0.001	0.586	0.609
0.002	0.765	0.668
0.003	0.848	0.699
0.005	1.052	0.823

**Table 4:** An evaluated results of twice the fraction of molecules as a function of the detuning for an adiabatic sweep through the resonance. The total density is equal to  $n = 10^{13} \text{ cm}^{-3}$ . The calculations have been done for two initial temperature  $T = 2T_0$  and  $4T_0$ .

$\frac{4\pi^2\hbar^6\delta}{g^4m^3}$	2 x Fraction of molecules	
	$T = 2T_0$	$T = 4T_0$
-0.0030	1.052	0.925
-0.0025	0.975	0.908
-0.0020	0.802	0.872
-0.0015	0.652	0.807
-0.0010	0.518	0.627
-0.0005	0.487	0.526
0.000	0.586	0.487
0.001	0.427	0.352
0.002	0.364	0.307
0.003	0.309	0.297
0.004	0.225	0.226

## CONCLUSION

From the above theoretical evaluations and investigations, we have come across the following conclusions.

- (1) At positive detuning, most of the atoms are free. One finds that the width in detuning of the cross-over regime is approximately equal to temperature and the result does not incorporate the effects of the nonzero lifetime of the molecules at positive detuning.
- (2) For negative detuning, one observes that the ideal gas result becomes equal to the exact result. This shows that for negative detuning, one is allowed to treat the gas as an ideal-gas mixture of atoms and dressed molecules.

## REFERENCES

1. A. J. Leggett. Rev. Mod. Phys (2001) **63**, 307
2. C. J. Pethick, H. Smith, “ Bose-Einstein condensation in dilute gases ( Cambridge University Press, Cambridge 2002)
3. L. P. Pitaevskii, S. Stringari, “ Bose-Einstein condensation (Oxford University Press, Oxford 2003)
4. V. Vuletic, A.J. Kerman, C. Chin and S. Chu, Phys. Rev. Lett. (PRL) (1999) **82**, 1409
5. H. Feshbach, Ann. Phys (1962) **19**, 287
6. R. A. Duine and H. T. C. Stoof, Phys Rev (2002) **A65**, 013603
7. S. K. Adhikari, Phys. Lett. (2002) **A296**, 145
8. H. Saito and M. Ueda, Phys. Rev (2002) **A65**, 033624
9. L. Santos and G. V. Shlyapnikov, Phys. Rev. (2002) **A66**, 011602
10. R. A. Duine and H. T. C. Stoof, Phys. Rev. (2003) **A68**, 013421
11. J. W. S. Negele and H. Orland, “ Quantum Many-Particle System (Addison-Weasley, New York, 1988)
12. S J. J. M. F. Kokkelmans, G. V. Shlyapnikov and G. Salomon, Phys. Rev. (2004) **A69**, 031423
13. P. Nozieres and S. Schmitt-Rink, J. Low Temp. Phys (JLTP) (1985) **59**, 195
14. R. A. Dunie and H. T. C. Stoof, New. J. Phys. (2003) **5**, 59
15. R. A. Duine and H. T. C. Stoof, Physics Reports (2004) **396**, 115
16. S. Sachdev, Phys. Rev. Lett. (PRL) (2003) **90**, 033201
17. B. J. Powell and R. H. McKenziel, Phys. Rev. Lett. (PRL) (2005) **94**, 047004
18. J. Geremia, J. Stockhom and H. Mubachi, Phys. Rev.(2006) **A73**, 042112
19. K. D.Nelson, Xiao Li and D. S. Weiss, Nature Physics (2007) **3**, 566
20. W. S. Bakr, J. I. Gillen, A. Peng, S. Folling and M. Greiner, Nature (2009) **462**, 7269
21. Y. Lin, S. Jung, S. Maxwell L. Turner , E. Tiesinga and P. Lett. (Phys. Rev. Lett.(PRL)(2009) **102**, 125301
22. D. V. Vasilyev, I. V. Sokolov and E. S. Polzik, Phys. Rev.(2010) **A 81**, 020302R
23. E. Zuethan, A. Gordeeka and A. Sorensen, Phys. Rev.(2011) **A84**, 043838
24. K. Hammer, A. S. Sorensen and F. S. Polzik, Rev. Mod. Phys.(2011) **84**, 1041

**Corresponding author: Indu Kumari Tiwari;** c/o Shree Anant Kumar  
Tiwari, Village-Dhawari, P.o-Bara Khayawani, Dist-Bhojpur -802166 (Bihar)