

Journal of Chemical, Biological and Physical Sciences

An International Peer Review E-3 Journal of Sciences

Available online at www.jcbpsc.org

Section C: Physical science



CODEN (USA): JCBPAT

Research Article

Electrical Conductivity and Dielectric Constant as Predictors of Chemical Properties and available Nutrients in the Soil

P. R. Chaudhari* and D. V. Ahire

Microwave Research Laboratory, Z. B. Patil College, Dhule-424002, India

Received: 20 February 2013; **Revised:** 7 March; **Accepted:** 9 March 2013

Abstract: The aim of this study was to evaluate chemical parameters and status of available total primary macronutrients (N+ P+ K), total secondary macronutrients (Ca+Mg) and total micronutrients (Cu+ Fe+ Mn+ Zn) in the soil from its electrical conductivity and dielectric constant. We have collected 8 samples (depth 0 - 15 cm) from different locations in North Maharashtra region. The soils were categorized as loamy sand, sandy loam and clay loam. The soils were analyzed for the chemical properties like organic carbon (OC) content, CaCO_3 , pH and for status of available nutrients. A pH and Electrical Conductivity of soil samples were measured by Soil Testing Kit. The dielectric constant was measured at C - band microwave frequency 4.5 GHz by using waveguide cell method. The statistical analysis of result shows high degree positive correlation of electrical conductivity and dielectric constant of soil samples with organic carbon (OC) content and available nutrients in soil samples while negative correlation with CaCO_3 and pH of soil samples. Regression equations are obtained to evaluate chemical properties and status of available nutrients from electrical conductivity and dielectric constant of soil.

Keywords: Electrical conductivity, dielectric constant, nutrients, microwave frequency.

INTRODUCTION

Soil is an important natural resource and plays a crucial role in maintaining environmental balance. Soil fertility is one of the important factors controlling yields of the crops. Soil characterization in relation to evaluation of fertility status of the soils of an area or region is an important aspect in context of sustainable agriculture production. Soil plays major role in determining the sustainable productivity of an agro-eco system. The sustainable productivity of a soil mainly depends upon its ability to supply essential nutrients to the growing plants. Proper nutrition is essential for satisfactory crop growth and production.

Soil has physico-chemical as well as electrical properties. Spatial and temporal variability of soil physical and chemical properties within a field are unavoidable. Color, texture, bulk density, nutrients and pH can be defined as its physico-chemical properties while dielectric constant, electrical conductivity and permeability are electrical properties.

Soil pH measurement is useful because it is a predictor of various chemical activities within the soil and hence can be used as a rough indicator of nutrients in the soil.

Laboratory soil test is usually time consuming and laborious. Since farmers cultivate annual crops more than once a year, this leads to delay in remedial action for the coming season. Hence, rapid measurement and monitoring of soil nutrient variability is needed to satisfy the precision farming requirements. These problems can be solved by measuring electrical conductivity (EC) of soil. Robert I. Papendick and James F. Parr¹ reported that soil test can help in determining the status of available nutrients to develop fertilizer recommendations for optimum crop production. M. Kumar and A.L. Babel² showed that the amount and availability of nutrients is controlled by physico-chemical and electrical properties of soil. The particle shape of soils, soil moisture contents and presence of chemical properties influence electrical conductivity of soil³. Electrical conductivity has generally been associated with determining soil salinity; however, EC also can serve as a measure of soluble nutrients⁴. In general, EC can be influenced by a number of different soil properties, including clay content and soil water content⁵.

The dielectric constant is a complex number and is a function of frequency. The real part of the dielectric constant of soil particles is between 3 and 6 depending on the orientation of the particles. Calla O.P.N. et al.⁶ have carried out extensive and systematic studies on the dielectric behavior of soils of Rajasthan and also from many parts of northern India.

In the present work, our aim is to measure electrical conductivity and dielectric constant of soil samples from North Maharashtra Region and use them to evaluate chemical parameters, status of available total primary macronutrients, total secondary macronutrients and total micronutrients in the soil.

MATERIAL AND METHODS

Soil samples (0-15cm) were collected from 8 sites covering North Maharashtra Region, keeping in view the physiographic characteristic in different cross sections of the area as well as variation in soil texture. Soils were completely air dried and passed through 2mm sieve and stored in properly labeled plastic bags for analysis. The sieved out fine particles are then oven dried to a temperature around 110⁰ C for several hours in order to completely remove any trace of moisture.


The samples were analyzed for the status of available nutrients by standard analytical methods from Govt. College of Agriculture, Dhule. The Chemical properties and nutrient concentrations of soil samples are represented in **Table1**.

Table-1: Nutrients and Chemical Properties of Soil Samples

Soil Parameters →	Total Primary Macro nutrients	Total Secondary Macro nutrients	Total Micro nutrients	OC	CaCO ₃	pH
Sample No. and Locations ↓	(N+P+K)	(Ca+Mg)	(Fe+Mn+ Zn+Cu)			
1. Amalner	864.9	247	22.9	0.65	4	7.2
2. Erandol	817.2	241	20.74	0.7	4.2	7.6
3. Shirpur	765.0	217	15.36	0.42	4.9	8
4. Dhule	693.4	216	16.20	0.5	5.5	8
5. Shahada	714.3	155	16.24	0.43	5.8	7.9
6. Kalwan	664.8	166	13.44	0.3	5.7	8.4
7. Satana	678.0	143	09.41	0.4	3.9	7.8
8. Nandurbar	581.2	116	08.87	0.23	6.4	8.25

EXPERIMENTAL

Measurement of Electrical Conductivity and pH of Soil Samples: A pH and Electrical Conductivity of soil samples were measured by Soil Testing Kit Model 161E. A 20 gm of collected soil was weighed out into a 150 ml plastic jar and 100 ml distill water was added to it. Lid of jar was packed tightly and stirred continuously for 5 minutes. Then it was kept overnight and stirred again. Allowed to set for 15 minutes and strained sample into clean measuring cup. A pH and Electrical conductivity readings were taken.

**Fig.1: Photograph of C-band microwave bench**

Measurement of Dielectric Constant of dry Soil Samples: The waveguide cell method⁶⁻⁸ is used to determine the dielectric properties of the dry soil samples. An automated C-band microwave set-up in the TE₁₀ mode with Gunn source operating at frequency 4.5 GHz, PC-based slotted line control and data acquisition system is used for this purpose. The solid dielectric cell with soil sample is connected to the opposite end of the source. The signal generated from the microwave source is allowed to incident on the soil sample. The sample reflects part of the incident signal from its front surface. The reflected wave combined with incident wave to give a standing wave pattern. These standing wave patterns are then used in determining the values of shift in minima resulted due to before and after inserting the sample. Experiments were performed at room temperatures ranged between 25°-35° C. Other details of dielectric constant measurement with C-band microwave bench set-up can also be seen from **Fig.1**

Dielectric constant (ϵ')

The dielectric constant ϵ' of the soils is determined from equation 1.

$$\epsilon' = \frac{g_{\epsilon} + \left(\lambda_{gs} / 2a \right)^2}{1 + \left(\lambda_{gs} / 2a \right)^2} \quad \dots 1$$

Where, a = Inner width of rectangular waveguide

λ_{gs} = wavelength in the air-filled guide, g_{ϵ} = real part of the admittance

The values of electrical conductivity and dielectric constant are listed in **Table 2**.

Table 2: Electrical Conductivity (EC) and Dielectric (DC) of Soil Samples

Sample No.→ Property ↓	1	2	3	4	5	6	7	8
EC(dSm ⁻¹)	0.33	0.30	0.26	0.17	0.20	0.18	0.15	0.10
DC	3.85	3.55	3.7	3.45	3.5	3.48	3.3	3.1

STATISTICAL ANALYSIS

The correlation coefficient 'r' of dielectric constant with soil parameters and nutrient concentration is determined using equation 2.

$$r = \frac{n \sum xy - (\sum x)(\sum y)}{\sqrt{n(\sum x^2) - (\sum x)^2} \sqrt{n(\sum y^2) - (\sum y)^2}} \quad \dots 2$$

Where n is the number of pairs of data (x, y).

The correlation coefficients (r) of dielectric constant with nutrient concentration and chemical parameters of soil samples and corresponding regression equations are given in **Table 3**.

Table 3: Correlation Coefficients and Regression Equations

Soil Nutrients and Parameter (y)	Correlation Coefficient 'r'		Regression Equations	
	with EC and Level of Significance	with DC and Level of Significance	EC	DC
(N+P+K)	0.9829 High Degree Positive	0.8888 High Degree Positive	$y = 1130 * EC + 483.5$	$y = 352.0 * DC - 509.7$
(Ca+Mg)	0.8849 Strong Positive	0.8567 Strong Positive	$y = 549.6 * EC + 71.50$	$y = 183.2 * DC - 453.8$
(Fe+Mn+Zn+Cu)	0.9160 High Degree Positive	0.8532 Strong Positive	$y = 57.17 * EC + 3.317$	$y = 18.34 * DC - 48.79$
pH	-0.7635 Strong Negative	-0.6008 Significant Negative	$y = -3.637 * EC + 8.662$	$y = -0.986 * DC + 11.34$
OC	0.8485 Strong Positive	0.6812 Significant Positive	$y = 1.729 * EC + 0.088$	$y = 0.478 * DC - 1.219$
CaCO ₃	-0.6465 Significant Negative	-0.4990 Significant Negative	$y = -7.727 * EC + 6.682$	$y = -2.054 * DC + 12.24$

RESULTS AND DISCUSSION

Soils were categorized as Clay, Silty Clay Loam, Clay Loam, Loamy, Sandy clay loam and Sandy Loam. All eight soil samples were found to be moderately calcareous (3.9 – 6.4 %) in nature. The pH (7.2 - 8.4) values and electrical conductivity values (0.10 – 0.33 dS/ m) indicated that all soil samples were found to be slightly to moderately alkaline and non saline in nature.

According to Methods Manual, Soil Testing in India⁹ the critical limits of Nitrogen (N), Phosphorus (P) and Potassium (K) for normal growth of plant were 280 kg/ha, 10 kg/ha and 108 kg/ha respectively. Consideration of these the available N (160 - 238 kg/ha) was low, P (6.2 - 16.9 kg/ha) of soils was ranging from low to high, while the available K (412 - 610 kg/ha) was very high. All soil samples were containing adequate amount of available Calcium (Ca) (98 - 200 meq/100 gm) and Magnesium (Mg) (18 – 49 meq/100 gm).

The critical limit of Fe is considered as 4.5 ppm for normal growth. Considering this limit only two soil samples 1 and 2 were containing sufficient amount (4.6 and 4.8 ppm) while the remaining soil samples were found to be deficient in available Fe (2.0 – 3.9 ppm).

The critical limit for available Mn is reported as 4.7 ppm. All soil samples were appeared to be high in available Mn (5.6 - 15 ppm).

The critical limit suggested for Zn is 0.5 to 1.00 ppm, according to which all soil samples are quite sufficient in available Zn (0.52 – 1.34 ppm) except sample 8 (0.32ppm).

Considering 0.66 ppm as critical limit of Cu for normal growth of plant, it may be inferred that all soil samples were contain adequate amount of available Cu (0.65 - 2.1 ppm).

For statistical analysis instead of individual content of nutrients we considered the total primary macronutrients (N+P+K) ranging from 581.2 to 864.9 kg/ha, total secondary macronutrients (Ca+Mg) ranging from 116 to 247 kg/ha and total micronutrients (Fe+Mn+Zn+Cu) ranging from 8.87 to 22.9 kg/ha.

Relationship of Electrical Conductivity and Dielectric Constant with total Macronutrient status of soil: High degree Positive correlation of electrical conductivity was found with available total primary macronutrients (N+ P+ K) ($r = 0.9829$) and available secondary macronutrients (Ca+ Mg) ($r = 0.8849$) of soil.

As that of electrical conductivity there was Strong Positive correlation of dielectric constant with available total primary macronutrients (N+ P+ K) ($r = 0.8888$) and available secondary macronutrients (Ca+ Mg) ($r = 0.8567$) of soil.

Relationship of Electrical Conductivity and Dielectric Constant with total Micronutrient status of soil: High degree Positive correlation was observed between electrical conductivity and total micronutrients (Fe +Mn +Zn +Cu) ($r = 0.9160$) of soil. Similarly there was Strong Positive correlation between dielectric constant and total micronutrients (Fe +Mn +Zn +Cu) ($r = 0.8532$) of soil.

Thus it was seen that electrical conductivity and dielectric constant of soil are positively correlated and very sensitive to total macronutrient and total micronutrient concentrations of soil.

Relationship of Electrical Conductivity and Dielectric Constant with pH, Organic Carbon and CaCO₃ of soil: Strong negative and strong positive correlations of electrical conductivity respectively with pH ($r = -0.7635$) and organic carbon ($r = 0.8485$) of soil were observed. While significant negative correlation was found between electrical conductivity and CaCO₃ ($r = -0.6465$) of soil.

On the other hand it was found that dielectric constant has significant negative correlation with pH ($r = -0.6008$) and CaCO₃ ($r = -0.4990$) while significant positive correlation with organic carbon ($r = 0.6812$) of soil.

Similar findings about relationship between available micronutrients, pH and electrical conductivity of soil were reported by M.Kumar and A.L. Babel² and R.H. Heiniger et al.¹⁰

CONCLUSIONS

1. High degree positive correlation of electrical conductivity and dielectric constant of soil samples with available total primary macronutrients (N+P+K), total secondary macronutrients (Ca+Mg) and total micronutrients (Fe+Mn+Zn+Cu) is obtained.
2. High degree positive correlation of electrical conductivity and dielectric constant of soil samples with organic carbon content of soil is observed.
3. High degree negative correlation of electrical conductivity and dielectric constant of soil samples with pH and significant negative with CaCO₃ content of soil is obtained.
4. The values of chemical parameters and status of available nutrients can be obtained from measured electrical conductivity and/or dielectric constant of soil with the use of regression equations in Table 3.

5. As compared to dielectric constant, electrical conductivity has strong significant correlations with total nutrient concentrations and chemical parameters of soil samples.

This study helps in determining the values of different chemical parameters and the nutrient concentrations of soil from North Maharashtra region by using known electrical conductivity and/or dielectric constant of soil.

REFERENCES

1. Robert I. Papendick and James F. Parr, *Amer. Journal of Alternative Agric.*, 1992, 7, 2.
2. M. Kumar and A.L. Babel, *India Journal of Agricultural Science*, 2011, 3, 97-106.
3. Z. Chik and T. Islam, *Int. J. Electrochemical Sci.*, 2011, 6, 6733- 6740.
4. J.L. Smith and J.W. Doran, Methods for Assessing Soil Quality, *Soil Sci. Soc. Am. Special Publication*, 1996, SSSA, Madison, WI.
5. R.G. Kachanoski, E. DeJong, and I.J. Van-Wesenbeeck, *Can. J. Soil Sci*, 1990, 70, 537-541.
6. O.P N Calla *Indian J. Radio Space Phys.* 1999, 28, 109-112.
7. P. R. Chaudhari, D.V. Ahire and V.D. Ahire, *Journal of Chemical, Biological and Physical Sciences*, 2012, 2, 3, 1493-1500.
8. D. H. Gadani and, A. D. Vyas, *Indian J. of Radio and Space Physics*, 2008, 37, 221- 229.
9. *Methods Manual*, Soil Testing in India, Department of Agriculture and Cooperation, Ministry of Agriculture, Government of India ,2011.
10. R.W. Heiniger, R.G. McBride and D.E. Clay, *Agron. Journal*, 2003, (95), 508-519.

Corresponding author: P. R. Chaudhari, Microwave Research Laboratory, Z. B. Patil College, Dhule-424002, India